Method of using 4-anilinoquinazoline derivatives as analgesic and anti- inflammatory agents	
Patent Number:	US4322420
Publication date:	1982-03-30
Inventor(s):	KOBAYASHI SHINSAKU; KAMOSHITA KATSUO; NAGAI SHIGEKI; HONDA TAKEO; ODA KIROKU; FUJII KATSUTOSHI; KOBAYASHI TAKASHI; KOJIMA MIKIO
Applicant(s)::	SANKYO CO
Requested Patent:	US4322420
Application Number:	US19790074343 19790911
Priority Number (s):	JP19780111484 19780911
IPC Classification:	A61K31/505 ; C07D239/94
EC Classification:	C07D239/94
Equivalents:	☐ <u>BE878723</u> , CA1151168, ☐ <u>CH642361</u> , ☐ <u>DE2936705</u> , DK377079, ☐ <u>ES8100276</u> , ☐ <u>FR2435248</u> , ☐ <u>GB2033894</u> , ☐ <u>IT1119933</u> , JP1400162C, ☐ <u>JP55038325</u> , JP62006546B, ☐ <u>NL7906761</u> , ☐ <u>SE446337</u> , SE7907493, ☐ <u>US4464375</u>
Abstract	
A method of using 4-anilinoquinazoline derivatives of formula (I): (I) (in which: R1 represents a hydrogen atom, a halogen, atom, a trifluoromethyl group or a nitro group; R2 represents a hydrogen atom, a C1-C4 alkyl group, an alkoxy group or a halogen atom; and R3 represents a hydrogen atom or a C1-C4 alkyl group) and pharmaceutically acceptable salts thereof are, provided that R1 does not represents a hydrogen atom or a chlorine atom in the 6- position when R2 and R3 both represent hydrogen atoms, as an analgesic and anti-inflammatory in the treatment of mammals. These compounds can be prepared by heating the appropriate 4-haloquinazoline with an appropriate aniline or aniline derivative. Data supplied from the esp@cenet database - I2	

- Please see exhibit 6 -